

The Deuterium Abundance in the Local Interstellar Medium

Tijana Prodanović¹

*Department of Physics, University of Novi Sad
Trg Dositeja Obradovića 4, 21000 Novi Sad, Serbia*

Gary Steigman²

*Department of Physics and Astronomy, Ohio State University
191 W. Woodruff Ave., Columbus OH 43210-1117, USA*

Brian D. Fields³

*Department of Astronomy, University of Illinois
1002 W. Green St., Urbana IL 61801, USA*

ABSTRACT

As the Galaxy evolves, the abundance of deuterium in the interstellar medium (ISM) decreases from its primordial value: deuterium is “astrated”. The deuterium astration factor, f_D , the ratio of the primordial D abundance (the D to H ratio by number) to the ISM D abundance, is determined by the competition between stellar destruction and infall, providing a constraint on models of the chemical evolution of the Galaxy. Although conventional wisdom suggests that the local ISM (i.e., within $\sim 1 - 2$ kpc of the Sun) should be well mixed and homogenized on timescales short compared to the chemical evolution timescale, the data reveal gas phase variations in the deuterium, iron, and other metal abundances as large as factors of $\sim 4 - 5$ or more, complicating the estimate of the “true” ISM D abundance and of the deuterium astration factor. Here, assuming that the variations in the observationally inferred ISM D abundances result entirely from the depletion of D onto dust, rather than from unmixed accretion of nearly primordial material, a model-independent, Bayesian approach is used to determine the undepleted abundance of deuterium in the ISM (or, a lower limit to it). We find the best estimate for the undepleted, ISM deuterium abundance to be $(D/H)_{\text{ISM}} \geq (2.0 \pm 0.1) \times 10^{-5}$. This result is used to

¹E-mail: prodanvc@df.uns.ac.rs

²E-mail: steigman@mps.ohio-state.edu

³E-mail: bdfields@illinois.edu

provide an estimate of (or, an upper bound to) the deuterium astration factor, $f_D \equiv (D/H)_P / (D/H)_{ISM} \leq 1.4 \pm 0.1$.

1. Introduction

Deuterium is created in an astrophysically interesting abundance only during big bang nucleosynthesis (BBN) (Boesgaard & Steigman 1985; Steigman 2007), after which, in the post-BBN Universe, its abundance ($y_D \equiv 10^5(D/H)$) decreases monotonically due to the processing of gas through succeeding generations of stars where deuterium is completely destroyed (Epstein et al. 1976; Prodanović & Fields 2003). Consequently, deuterium plays a special role in cosmology, nuclear astrophysics, and in Galactic chemical evolution (Yang et al. 1984; Boesgaard & Steigman 1985; Steigman & Tosi 1992, 1995; Vangioni-Flam et al. 1994; Tosi 1996; Schramm & Turner 1998; Romano et al. 2006; Steigman et al. 2007; Steigman 2007; Prodanović & Fields 2008). Its relatively simple Galactic evolution permits us to use deuterium to determine the fraction of interstellar gas that has been processed through stars (Steigman & Tosi 1992, 1995). By comparing the primordial and Galactic deuterium abundances one can learn about and discriminate among different Galactic chemical evolution models (Vangioni-Flam et al. 1994; Tosi 1996; Romano et al. 2006; Steigman et al. 2007; Prodanović & Fields 2008). Together with non-BBN constraints on the baryon (nucleon) density from observations of the cosmic microwave background (Spergel et al. 2007; Dunkley et al. 2009; Komatsu et al. 2009, 2010), the primordial deuterium abundance, y_{DP} , is predicted by BBN (Cyburt et al. 2003; Coc et al. 2004; Steigman 2007; Simha & Steigman 2008; Cyburt et al. 2008). The predicted abundance is in excellent agreement with observations of deuterium in high-redshift, low-metallicity QSO absorption line systems (QSOALS) (O’Meara et al. 2006; Pettini et al. 2008).

Since deuterium is destroyed in the Galaxy as gas is cycled through stars, $(D/H)_{ISM} \leq (D/H)_P$. However, all successful chemical evolution models require infall to the disk of the Galaxy of unprocessed (or, nearly unprocessed) gas (Tosi 1996; Prodanović & Fields 2008), and such deuterium-rich (and metal-poor) gas would raise the ISM D/H ratio closer to the primordial value. As a result, comparison of the primordial and ISM deuterium abundances provides a constraint on infall and, therefore, on chemical evolution models. Observations over the past decade and more of the deuterium abundance in the relatively local interstellar medium (ISM) reveal an unexpectedly large scatter in D/H (Jenkins et al. 1999; Sonneborn et al. 2000; Hébrard et al. 2002; Hoopes et al. 2003), challenging the conventional wisdom of a well mixed ISM. For example, as shown in Figure 1, the absorption line measurements from the *Far Ultraviolet Spectroscopic Explorer* (FUSE) reveal variations




Fig. 1.— The logs of the deuterium abundances versus the logs of the H I column densities [cm^{-2}] for the 49 FUSE LOS (see the text). The filled symbols are for the 41 LOS which have iron abundance data, while the open symbols are for the 8 LOS which lack iron abundances. The squares (blue) are for the LOS within the Local Bubble (LB) and the circles (red) are for the non-LB (nLB) LOS. The solid line is at the mean D abundance for the LB LOS ($\log y_{\text{D,LB}} = 0.19$); the dashed line is its extension to the nLB LOS.

of a factor of ~ 4 ($0.5 \lesssim y_{\text{D,ISM}} \equiv 10^5 (\text{D}/\text{H})_{\text{ISM}} \lesssim 2.2$) in the *gas-phase* D/H ratios over lines of sight (LOS) to background stars within $\sim 1 - 2$ kpc of the Sun. Moreover, the variations in the observed, gas phase D/H abundances are found to correlate *positively* with the abundances of refractory elements such as Ti (Prochaska et al. 2005; Lallement et al. 2008), and a similar correlation is also found between D and other metals such as Fe and O (Linsky et al. 2006; Steigman et al. 2007). However, this positive correlation between the abundances of D and the metals is opposite to the trend expected from stellar nucleosynthesis (D decreasing as the metals increase). Motivated by the observed correlations and by the very large spread in the D/H ratios inferred from the FUSE and other data sets, it has been proposed that the large variations in *gas-phase* D/H may be due to the depletion of gas phase deuterium onto dust grains (Jura 1982; Draine 2004, 2006). If this is the case, then the FUSE (and other) gas-phase absorption-line measurements reveal that depletion has not been well mixed (homogenized) in the ISM and, the data may only provide a *lower* limit to the *true* (undepleted) ISM deuterium abundance and, therefore, only an *upper* limit to the deuterium astration factor, $f_{\text{D}} \equiv y_{\text{DP}}/y_{\text{D,ISM}}$.

In Figure 1 the logs of the deuterium abundances ($\log y_{\text{D}} \equiv 5 + \log N(\text{D I}) - \log N(\text{H I})$) are shown as a function of the logs of the H I column densities for the 49 LOS with D I column

densities from Table 2 of Linsky et al. (2006), supplemented by data from Oliveira & Hébrard (2006) and Dupuis et al. (2009). Of these 49 LOS, 41 have iron abundance measurements (filled symbols); 21 of the 49 LOS are within the Local Bubble (LB; see Linsky et al. (2006) for a discussion of the LB); the remaining 28 non-Local Bubble (nLB) LOS are toward stars beyond the LB (see §2.1).

The unexpectedly large spread among the observationally inferred ISM D abundances complicates any estimate of $y_{\text{D,ISM}}$. Recognizing this point, Linsky et al. (2006) chose for their estimate of a *lower bound* to the true (undepleted) ISM deuterium abundance the mean of the five *highest* D/H ratios finding $y_{\text{D,ISM}} \geq 2.17 \pm 0.17$ or, when including corrections (see the discussion in §2.1) for N(H I) and N(D I) for those lines of sight outside of the Local Bubble, $y_{\text{D,ISM}} \geq 2.31 \pm 0.24$. These estimates are quite close to the *lower bound* to the primordial abundance estimated from the QSOALS, $y_{\text{DP}} = 2.82^{+0.20}_{-0.19}$ (Pettini et al. 2008)¹, suggesting a small upper bound to the D astration factor, $f_{\text{D}} \leq 1.30^{+0.14}_{-0.13}$ or, an even smaller value, $f_{\text{D}} \leq 1.22 \pm 0.15$ for the LB-corrected, nLB deuterium abundances. To account for such a high ISM deuterium abundance and such a small D astration factor, a very high infall rate of pristine material would be needed, challenging many Galactic chemical evolution models (Romano et al. 2006; Prodanović & Fields 2008). By limiting themselves to the five highest D abundances, Linsky et al. (2006) ignore the lower deuterium abundances along many more LOS which are consistent with them within the errors, potentially biasing their estimates of $y_{\text{D,ISM}}$ and of f_{D} .

In an attempt to address this issue, Steigman et al. (2007) (SRT), used the 18 highest D/H ratios from the FUSE data (see Table 3 of Linsky et al. (2006)), finding an ISM D abundance of $y_{\text{D,ISM}} = 1.88 \pm 0.11$, corresponding to a D-astration factor $f_{\text{D}} \leq 1.50^{+0.14}_{-0.13}$, consistent with at least some of the otherwise successful chemical evolution models discussed in SRT. In fact, the data in Table 2 of Linsky et al. (2006) reveal that there are 19 LOS

¹ Since this estimate of y_{DP} relies on only 7 high redshift, low metallicity LOS and, since the dispersion in deuterium abundances among them is unexpectedly large, some prefer to adopt a so-called “WMAP D abundance”. WMAP does not observe deuterium. Rather, the WMAP-determined baryon density parameter may be used in a BBN code to *predict* the relic D abundance. If the Komatsu et al. (2010) estimate of the baryon density is adopted, the BBN predicted primordial D abundance is $y_{\text{DP}} = 2.5 \pm 0.1$. However, the WMAP collaboration also provides an estimate of the effective number of neutrinos, (Komatsu et al. 2010) which, when used along with their baryon density estimate in a BBN code, leads to a *different* predicted primordial D abundance $y_{\text{DP}} = 3.0 \pm 0.4$. The difference between these two predictions reflects the difference between the standard model effective number of neutrinos expected and the WMAP observed value, which is within $\sim 1.5\sigma$ of expectations. So, which, if either, “WMAP D abundance” is preferred? Here, we compare observations to observations, not to model-dependent predictions and, we adopt the Pettini et al. (2008) value for y_{DP} .

with central values of $\log y_D \geq 0.20$. The weighted mean (along with the error in the mean) for these 19 D abundances is $\log y_{D,19} = 0.26 \pm 0.01$, corresponding to $y_{D,19} = 1.8 \pm 0.1$. For these 19 LOS the reduced $\chi^2 = 0.85$, confirms that the weighted mean provides a good description of their D abundances. As more LOS with lower D abundances are added, the weighted mean decreases, but the reduced χ^2 increases, so that $y_{D,ISM} \gtrsim 1.8 \pm 0.1$ is likely a robust *lower* bound to the ISM D abundance and, $f_D \lesssim 1.5 \pm 0.1$ ($\log f_D \lesssim 0.19 \pm 0.03$) is a robust *upper* bound to the deuterium astration factor.

Surely, there must be a better way to find a reliable estimate of the maximum value of the deuterium abundance in the local ISM while accounting for the observational errors in the individual D abundance determinations. In §2 we describe a Bayesian analysis designed to find the best estimate of the maximum, gas phase (undepleted), ISM deuterium abundance, $y_{D,max}$, from data with non-negligible errors and we apply it to the FUSE data set. On the assumption that the spread in the observed D abundances is the result of incompletely homogenized dust depletion, $y_{D,ISM} \geq y_{D,max}$ and $f_D \equiv y_{DP}/y_{D,ISM} \leq y_{DP}/y_{D,max}$. Our results are summarized and our conclusions presented in §3. As already noted, the dust depletion hypothesis suggests that there should be a correlation between the D and Fe abundances. Although this trend appears to be present at some level (Linsky et al. 2006), a deeper analysis suggests that the simplest interpretation of the depletion hypothesis may need to be modified (Steigman et al. 2007). For example, since D is likely attached to the mantles of dust grains (Tielens 1983), while Fe is mostly contained in the cores of the grains, D will be removed from grains more easily than Fe. Thus, although strong shocks may destroy grains completely, weak shocks may only remove D from grains while Fe might stay locked within their cores (Draine & Salpeter 1979). This effect then may explain the scatter observed in the relation between the D and Fe depletions. Thus, in a follow-up paper in preparation, the consistency of the depletion hypothesis with the FUSE data for D and Fe (and O) is investigated, and some possible modifications to its simplest version are explored (Steigman & Prodanović 2010).

2. A Bayesian Analysis Of The ISM D Abundances

To avoid imposing any prior prejudice on which LOS should be included and which excluded in our analysis or, on how the observed D abundances may, or may not correlate with iron (or other metals), we adopt a statistical, model-independent method for determining the undepleted Galactic deuterium abundance. Our approach follows closely the model-independent Bayesian analysis developed by Hogan et al. (1997) to determine the primordial helium abundance. It is useful to recall the problem Hogan et al. (1997) addressed and how

they solved it. Helium-4 is produced abundantly during BBN and in the post-BBN Universe its abundance is enhanced by stellar produced ^4He . As a result, the ^4He mass fraction, Y_P , is expected to increase from its primordial value, Y_P , along with the metallicity, Z . Extrapolation of the observed Y versus Z relation to zero metallicity results in an estimate of $Y_{\min} = Y_P$. The problem for ^4He is that the form of the Y_P versus Z relation is a priori unknown and, may even differ from object to object (low metallicity, extragalactic H II regions). This, in combination with the errors in the observed values of Y_P and Z , complicates the derivation of Y_P from the data. The challenge confronting Hogan et al. (1997) was to identify $Y_{\min}$. The Bayesian approach they developed (Hogan et al. 1997) is described below. If, indeed, the observed spread in D abundances results from dust depletion, then their ^4He problem is entirely analogous to the one we confront in using the D abundance data, with its errors, to infer the *maximum*, gas phase D abundance.

Here, the FUSE set of Galactic ISM deuterium observations is analyzed assuming only that there exists a “true”, uniform, ISM deuterium abundance whose *gas phase* value may have been reduced by the depletion of D onto dust grains, with the amount of depletion possibly varying from LOS to LOS. The a priori unknown distribution of depletions is characterized in terms of a Bayesian probability distribution (a “prior”), and the data themselves are used to determine both the true ISM abundance $y_{\text{D,ISM}} = y_{\text{D,max}}$ or, a lower bound to it, $y_{\text{D,ISM}} \geq y_{\text{D,max}}$, as well as a measure of the amount of depletion, the depletion parameter $w \equiv y_{\text{D,max}} - y_{\text{D,min}}$. No prior assumptions about which LOS may have been affected by dust depletion are imposed and the entire data set is analyzed in an unbiased, Bayesian manner.

2.1. The Data

In our analysis we use the FUSE ISM deuterium abundance data for 46 LOS from Linsky et al. (2006), together with three, more recent measurements towards HD41161, HD53975 (Oliveira & Hébrard 2006) and REJ1738+665 (Dupuis et al. 2009). Of the 49 LOS with deuterium abundance measurements, 21 are within the LB (see Linsky et al. (2006) for a discussion of the LB); the remaining 28 are nLB LOS, towards stars beyond the LB. While the star 31 Com is more distant than all the LB stars, the LOS to it has a very low H I column density and an average H I volume density ($n(\text{H I}) \equiv N(\text{H I})/d$) much smaller than for all LB LOS. According to Piskunov et al. (1997) and Dring et al. (1997), the absorption feature toward 31 Com is at a velocity which is inconsistent with the LB and this LOS likely lies within the hot, ionized, Local Bubble (which may account for the “high” D and Fe abundances) in the direction of the north galactic pole. For a contrary point of view, see Redfield & Linsky (2008). In contrast to the assignment in Table 2 of Linsky et al. (2006), we include 31 Com

along with the nLB LOS (see Figure 1). However, this assignment has negligible impact on our quantitative results. Linsky et al. (2006) also list D/H ratios for the nLB LOS *corrected* by them for assumed *average* foreground LB D I and H I absorption. If velocity information were available it could be used to separate foreground absorption in the LB from that due to gas lying beyond the LB. In the absence of such data, Linsky et al. (2006) *assume* that the LB extends to $N(\text{H I})_{\text{LB}} = 10^{19.2}$ and multiply $N(\text{H I})_{\text{LB}}$ by an adopted average LB D/H ratio to find $N(\text{D I})_{\text{LB}}$. These two average column densities are subtracted from the D I and H I column densities observed for the individual nLB LOS and the ratio of the corrected column densities is used to find corrected nLB D/H ratios. Linsky et al. (2006) base their assumed LB H I column density on the Na I observations of Lallement et al. (2003), who find tentative evidence for a LB “wall” of cold dense gas with $N(\text{H I}) \sim 10^{19.5}$. This procedure can bias the “corrected” nLB deuterium abundances. For example, it enhances the D abundances along those nLB LOS where the observed D abundance exceeds the adopted LB D abundance and decreases the D abundances for those LOS where the observed D abundance is less than the adopted LB value. The magnitude of the correction increases with the value of the “average” H I column density adopted. In fact, the observed LB H I column and volume densities are distributed very inhomogeneously, varying by nearly a factor of 30 along different LB LOS. This results in a scatter plot for the column densities as a function of distance to the background star. Even for the subset of the most distant LB stars with $d \sim 70 - 80$ pc, $N(\text{H I})$ varies by a factor of ~ 16 , $10^{17.9} \lesssim N(\text{H I}) \lesssim 10^{19.1}$. The structure of the LB is very complex (Lallement et al. 2003) and the location of the LB “wall” varies from 65-150 pc. Thus, the value for the LB H I column density adopted by Linsky et al. (2006) may be an *overestimate*, leading to an *overestimate* for their inferred value of $y_{\text{D,max}}$, since a smaller choice for the average foreground LB H I column density will result in a smaller correction to $N(\text{D I})$ and a lower estimate of $y_{\text{D,max}}$. For these reasons, in our analysis, we use the uncorrected H I and D I column densities for all nLB LOS.

Following Linsky et al. (2006), we adopt as a working hypothesis that the large scatter in the observed, gas phase, ISM deuterium abundances is a reflection of the preferential depletion of deuterium (relative to hydrogen) onto dust grains. Therefore, it is assumed that in the ISM deuterium has a total LOS column density $N(\text{D})_{\text{ISM}} = N(\text{D})_{\text{gas}} + N(\text{D})_{\text{dust}}$, where the sum includes the observed, gas phase component, $N(\text{D})_{\text{gas}} = N(\text{D I})_{\text{OBS}}$, and an unobserved, dust-depleted component, $N(\text{D})_{\text{dust}}$. Along any LOS, $N(\text{D})_{\text{ISM}} \geq N(\text{D})_{\text{gas}}$. The data suggest, and we assume, that hydrogen is negligibly depleted onto dust and that the fraction of H tied up in molecules in the diffuse ISM probed by FUSE is usually negligible (when H_2 is observed, its column density is included in the budget for gas phase hydrogen), so that $N(\text{H})_{\text{ISM}} = N(\text{H})_{\text{gas}}$.

On the assumption of a uniform D abundance (gas plus dust) in the local ISM, the

dispersion among the observed gas phase D abundances, $y_{D,i}$, reflects the observational errors in $N(\text{D I})$ and $N(\text{H I})$, along with any spatial variations in D depletion. Since deuterium may be depleted along ALL local ISM LOS, $y_{D,i} \leq y_{D,\text{max}}$, where $y_{D,\text{max}}$ is the maximum gas phase (undepleted) deuterium abundance, consistent with the observational errors, and $y_{D,\text{ISM}} \geq y_{D,\text{max}}$. Within the local ISM there will be a maximum depletion so that $y_{D,i} \geq y_{D,\text{min}}$. Along the i th LOS, the observed, gas phase D abundance, $y_{D,i}$ ($y_{D,i} \equiv y_{D,i,\text{obs}}$), may differ from the maximum undepleted abundance by an amount w_i where, $w_i \equiv y_{D,\text{max}} - y_{D,i}$, and $0 \leq w_i \leq w \equiv y_{D,\text{max}} - y_{D,\text{min}}$. In the absence of observational errors, the observed D abundance along the i th LOS would lie between $y_{D,\text{min}}$ and $y_{D,\text{max}}$ and would reflect the difference between the maximum undepleted ISM abundance $y_{D,\text{max}}$, and the spatially varying value of w , $w_i = y_{D,\text{max}} - y_{D,i}$. Observational errors complicate the task of using the data, $\{y_{D,i}\}$, to identify $y_{D,\text{max}}$ (and $y_{D,\text{min}}$ or, w).

Since real data do have errors, the observationally inferred D abundance along the i th LOS, $y_{D,i}$, will differ from the true gas phase D abundance, $y_{D,i,\text{T}}$, by an amount δ_i ($y_{D,i} = y_{D,i,\text{T}} + \delta_i$). For an individual measurement the difference between the observed and true D (gas phase) abundances, δ_i , is unknown. If it is assumed that δ_i is a random variable drawn from a zero-mean Gaussian of width σ_i , then the probability distribution for $y_{D,i}$, given a true gas phase abundance $y_{D,i,\text{T}}$ along the i th LOS is

$$P(y_{D,i}|y_{D,i,\text{T}}) = \frac{1}{\sqrt{2\pi}\sigma_{i\pm}} e^{-(y_{D,i}-y_{D,i,\text{T}})^2/2\sigma_{i\pm}^2} \quad (1)$$

In eq. (1) we allow for asymmetrical measurement errors so that σ_{i+} corresponds to $y_{D,i} > y_{D,i,\text{T}}$ and σ_{i-} to $y_{D,i} < y_{D,i,\text{T}}$.

2.2. Bayesian Formalism

For a set of data with errors $\{y_{D,i}, \delta_i\}$, the Bayesian approach described by Hogan et al. (1997) enables a determination of two parameters, $y_{D,\text{max}}$ and w (or $y_{D,\text{min}}$). Along each LOS in the local ISM the true abundance is related to $y_{D,\text{max}}$ by $y_{D,i,\text{T}} = y_{D,\text{max}} - w_i$.

The likelihood of finding particular values of $y_{D,\text{max}}$ and of w (or $y_{D,\text{min}}$), $\mathcal{L}(y_{D,\text{max}}; w)$, from a sample of measurements $y_{D,i}$, with errors δ_i is,

$$\mathcal{L}(y_{D,\text{max}}; w) = \prod_i P(y_{D,i}|y_{D,\text{max}}; w) \quad (2)$$

$$\mathcal{L}(y_{D,\text{max}}; w) = \prod_i \int dy_{D,i,\text{T}} P(y_{D,i}|y_{D,i,\text{T}}) \times P(y_{D,i,\text{T}}|y_{D,\text{max}}; w) \quad (3)$$

where $P(y_{D,i}|y_{D,i,T})$ is the probability distribution from eq. (1), relating each observed abundance $y_{D,i}$ to its underlying true value $y_{D,i,T}$. $P(y_{D,i,T}|y_{D,max}; w)$ is the dust depletion probability distribution – the probability of finding the true, dust depleted, LOS deuterium abundance $y_{D,i,T}$, given the values of the maximum and minimum ISM deuterium abundances, $y_{D,max}$ and $y_{D,min} = y_{D,max} - w$ respectively. Our probability distribution, $P(y_{D,i,T}|y_{D,max}; w)$, determines the distribution of the w_i . To avoid any a priori model dependence and to limit ourselves to the fewest assumptions, following Hogan et al. (1997) we initially adopt the simplest form for this probability distribution – a *top-hat* function:

$$P(y_{D,i,T}|y_{D,max}; w) = \begin{cases} 1/w & , \quad y_{D,min} \leq y_{D,i,T} \leq y_{D,max} \\ 0 & , \quad \text{otherwise} \end{cases} \quad (4)$$

where $y_{D,min} = y_{D,max} - w$. The probability distribution is normalized to unity when integrated over $y_{D,i,T}$. Integrating over the range of possible deuterium abundances for N data points, we find the likelihood distribution for the maximum gas phase deuterium abundance in the local ISM, $y_{D,max}$, and for w ,

$$\mathcal{L}(y_{D,max}; w) = \prod_i^N \frac{1}{2w} \left[\text{erf} \left(\frac{y_{D,i} - y_{D,min}}{\sqrt{2}\sigma_{i+}} \right) - \text{erf} \left(\frac{y_{D,i} - y_{D,max}}{\sqrt{2}\sigma_{i-}} \right) \right] \quad (5)$$

Equation (5) is evaluated numerically for a range of $y_{D,max}$ and w (or $y_{D,min}$) values to find the combination of these two parameters which maximizes the likelihood. In this way the set of deuterium observations, including the errors, is used to find the most likely values of $y_{D,max}$ and the depletion parameter, w , avoiding any prior assumptions about which LOS may, or may not, be affected by depletion of deuterium onto dust or, about how the abundance of D may, or may not, be correlated with the abundance of iron or other metals.

In addition to adopting the top-hat depletion probability distribution, following Hogan et al. (1997) we also explore the consequences of choosing two other, asymmetric probability distributions:

$$P(y_{D,i,T}|y_{D,max}; w) = \begin{cases} 2(y_{D,i,T} - y_{D,min})/w^2 & , \quad y_{D,min} \leq y_{D,i,T} \leq y_{D,max} \\ 0 & , \quad \text{otherwise} \end{cases} \quad (6)$$

$$P(y_{D,i,T}|y_{D,max}; w) = \begin{cases} 2(y_{D,max} - y_{D,i,T})/w^2 & , \quad y_{D,min} \leq y_{D,i,T} \leq y_{D,max} \\ 0 & , \quad \text{otherwise} \end{cases} \quad (7)$$

where eq. (6) is a *positive bias* distribution, favoring smaller depletion of deuterium, while eq. (7) is a *negative bias* distribution, favoring larger D-depletion. At the Referee's suggestion we have also analyzed the effects of another probability distribution – a bimodal, *M-shaped*

bias distribution defined in eq. (8), which favors both low and high depletion of deuterium, while moderate depletion is disfavored. Finally, we also explored the complementary, *Λ-shaped* probability distribution presented in eq. (9). Both additional distributions are defined with respect to the midpoint $y_M = (y_{D,\max} + y_{D,\min})/2 = y_{D,\max} - w/2$.

$$P(y_{D,i,T}|y_{D,\max}; w) = \begin{cases} \frac{2(y_M - y_{D,i,T})}{w(y_M + w - y_{D,\max})} , & y_{D,\min} \leq y_{D,i,T} \leq y_M \\ \frac{2(y_M - y_{D,i,T})}{w(y_M - y_{D,\max})} , & y_M \leq y_{D,i,T} \leq y_{D,\max} \end{cases} \quad (8)$$

$$P(y_{D,i,T}|y_{D,\max}; w) = \begin{cases} \frac{2(y_{D,i,T} - y_{D,\max} + w)}{w(y_M + w - y_{D,\max})} , & y_{D,\min} \leq y_{D,i,T} \leq y_M \\ \frac{2(y_{D,i,T} - y_{D,\max})}{w(y_M - y_{D,\max})} , & y_M \leq y_{D,i,T} \leq y_{D,\max} \end{cases} \quad (9)$$

All probability distributions are normalized to unity when integrated over $y_{D,i,T}$.

Our Bayesian analysis determines two parameters, the best fit values of the maximum and minimum deuterium abundances, $y_{D,\max}$ and $y_{D,\min} = y_{D,\max} - w$, compatible with the errors in the data and the assumption that any spread in the observed abundances, above and beyond that expected from the errors, is real and reflects the variable ISM depletion of deuterium onto dust. Hogan et al. (1997) were mainly interested in the *minimum* ^4He abundance, $Y_{\min}$, and we are primarily interested in the *maximum* ISM D abundance, $y_{D,\max}$. The likelihood in the $\{y_{D,\max}, w\}$ plane is maximized to find the best estimate of the maximum undepleted D abundance, providing a lower limit to the ISM D abundance ($y_{D,\text{ISM}} \geq y_{D,\max}$). We note that in the absence of observational errors, the depletion parameter, w , is restricted to $0 \leq w \leq y_{D,\max}$, since $y_{D,\max} \geq y_{D,\min}$. In Figures 2 – 4, w starts at zero and the dashed lines show the boundary, $w = y_{D,\max}$.

2.3. Results

As may be seen from Figure 1, the distributions of the observed D abundances for the LB and nLB LOS are very different. The LB y_D values show little gas phase variation from LOS to LOS, in contrast to the nLB D/H ratios, whose variation accounts for most of the factor ~ 4 spread in the observed ISM D abundances. Indeed, all 21 LB D abundances are consistent with no variation (within the observational errors) around a weighted mean of $\log(y_{D,\text{LB}}) = 0.19$, corresponding to $y_{D,\text{LB}} = 1.5$. Our Bayesian analysis for all five probability distributions is in agreement with this and yields $w \approx 0$ and $y_{D,\max} \approx y_{D,\min} \approx 1.5$. Our results for three distributions (top-hat, positive-bias, negative-bias), shown in Figure 2, illustrate this result. A *lower* bound to the deuterium astration factor may be inferred from the *upper* bound to the LB deuterium abundances, $y_{D,\max} = y_{D,\text{LB}}$, and the estimate of y_{DP} from Pettini et al.

(2008). For central values of $\log(y_{\text{D, LB}}) \geq 0.19$ and $\log(y_{\text{DP}}) = 0.45$ (Pettini et al. 2008), $\log(f_{\text{D, LB}}) \leq 0.26$, corresponding to $f_{\text{D, LB}} \leq 1.8$ ², consistent with the successful Galactic chemical evolution models identified in SRT. This value is also consistent with a wide range of chemical evolution models discussed in Prodanović & Fields (2008) with both low and high infall rates of nearly pristine gas, as well as with a variety of initial mass functions.

In contrast to the LB, the nLB LOS do show large variations among the observed gas phase D abundances (see Figure 1). It is thus expected that the Bayesian analysis of this data subset will find evidence for $w > 0$ at a statistically significant level. This indeed is seen in Figures 4 (top-hat) and 5 (M-shaped bias) where our results are shown separately for the data of the LB (top panel), the nLB (middle panel) LOS, along with those for all 49 LOS (bottom panel). For the 28 nLB LOS the best fit values, from a top-hat distribution, are $y_{\text{D, max}} = 2.1$ and $w = 1.5$. The top-hat Bayesian analysis for the complete FUSE data set (Figure 4, bottom panel) finds maximum likelihood values of $y_{\text{D, max}} = 2.0$ and $w = 1.3$, corresponding to $y_{\text{D, min}} = 0.7$. The non-zero depletion parameter found for all 49 LOS is driven by the large variations in y_{D} for the nLB LOS. This value of $y_{\text{D, max}}$ corresponds to $f_{\text{D}} \leq 1.4$, which, within the errors in the measurements of y_{DP} and $y_{\text{D, max}}$, is marginally consistent with the fiducial chemical evolution model discussed in SRT and is consistent with a range of models discussed in Prodanović & Fields (2008), where even larger infall rates are required for some initial mass functions. On the other hand, when the complete FUSE data set is analyzed with the M-shaped prior suggested by the referee (Figure 5, bottom panel), the maximum likelihood is obtained for $y_{\text{D, max}} = 1.8$ and $w = 1.0$, corresponding to $y_{\text{D, min}} = 0.8$ and $f_{\text{D}} \leq 1.6$. This result is consistent with the fiducial model adopted in SRT, but not with some of the other Galactic chemical evolution models discussed by them (Steigman et al. 2007). Similar to the nLB case, this result is also consistent with some of the models explored in Prodanović & Fields (2008), where the observations allow for both high and low infall rates depending on the choice of the initial mass function. However, the more recent initial mass functions are only consistent with larger infall rates.

²Significant figures and round off: The H I and D I column densities listed in Table 2 of Linsky et al. (2006) are typically given to two significant figures (the integers in front of the decimal place simply count the powers of ten). These column densities are used to find $\log(y_{\text{D}}) = 5 + \log N(\text{D I}) - \log N(\text{H I})$, where the errors in the logs of the column densities are added in quadrature. As a result, the values of $\log(y_{\text{D}})$ (and their errors) are only known to **two** significant figures and, so too are the values of y_{D} (and their errors) derived from them. However, in Table 3 of Linsky et al. (2006) the values of y_{D} are given to **three** (or more) significant figures. For our Bayesian analysis we use the data in Table 3 of Linsky et al. (2006), but the results presented here are generally rounded to two significant figures. As an example, if we had first rounded the individual values of $y_{\text{D, LB}}$ and y_{DP} to two significant figures, we would have found $y_{\text{D, LB}} = 1.5$ and $y_{\text{DP}} = 2.8$, leading to $f_{\text{D}} \leq y_{\text{DP}}/y_{\text{D, LB}} = 1.9$, in contrast to the slightly different value quoted here, which follows from $\log(f_{\text{D}}) \leq \log(y_{\text{DP}}) - \log(y_{\text{D, LB}}) = 0.26$ and $f_{\text{D}} = 10^{0.26} = 1.8$.




Fig. 2.— Likelihood contours (68%, 95%, 99%) in the $w - y_{D,\max}$ plane for the 21 LB LOS for the top-hat (top panel), the positive-bias (middle panel), and the negative-bias probability distribution (bottom panel), where $w \equiv y_{D,\max} - y_{D,\min}$ is the depletion parameter. The best fit values – for all three probability distributions – are at $y_{D,\max} = 1.5$ and $w = 0$ ($y_{D,\min} = y_{D,\max}$), rounded off to two significant figures. The dashed line, $w = y_{D,\max}$, represents the bound above which the results, in the absence of errors, are unphysical since $y_{D,\min} = y_{D,\max} - w$ becomes negative.

As anticipated from Figure 1, the bottom panels of Figures 4 and 5 confirm that the variation among the gas phase D abundances for all 49 FUSE LOS is too large to be accounted for by the observational errors ($w \neq 0$ at much greater than 99% confidence), for all adopted probability distributions. Indeed, *e.g.*, for the top-hat probability distribution, the D abundances for all 49 LOS span a range of nearly a factor of three, from $y_{D,\max} = 2.0$, down to $y_{D,\min} \equiv y_{D,\max} - w = 0.7$.

The effects of the choices of the priors on the likelihood distributions for $y_{D,\max}$ and w are shown for the complete FUSE data set on Figure 3 and on the bottom panel of Figure 5. While the results for the top-hat and positive bias distributions are quite similar, the negative-bias distribution is noticeably different. The best fit parameters $y_{D,\max}$, w and resulting astration factor f_D are summarized in Table 1. Also presented in the table is, $\Delta \ln \mathcal{L}_{\max}$, the difference between the logarithms of the largest maximum likeli-




Fig. 3.— Likelihood contours (68%, 95%, 99%) in the $w - y_{D,max}$ plane for all 49 LOS for the top-hat (top panel), the positive-bias (middle panel), and the negative-bias probability distribution (bottom panel). The dashed line is $w = y_{D,max}$; see Figure 2.

hood and a maximum likelihood of a given probability distribution. While the M-shaped probability distribution has the largest maximum likelihood value, so that $\Delta \ln \mathcal{L}_{max} \equiv \ln \mathcal{L}_{max,M} - \ln \mathcal{L}_{max,bias}$, this distribution is closely followed by the top-hat and positive-bias priors, suggesting that $1.8 \lesssim y_{D,max} \lesssim 2.0$. The negative-bias prior, favoring large depletion, yields the poorest fit to the data. As may be seen from Figure 1, the large errors in the data mask which of our priors might provide the best fit. Because the top-hat distribution requires the simplest assumption (no preference), and since its maximum likelihood is similar to that of the M-prior, we to adopt $y_{D,max} = 2.0 \pm 0.1$ as our best estimate for the maximum of the gas phase, ISM D abundances.

3. Summary and Conclusions

In the decades prior to the current era of precision cosmology the primordial abundance of deuterium provided the only quantitative cosmological baryometer (Boesgaard & Steigman 1985). Although, at present, the deuterium abundance is only measured along seven, high-




Fig. 4.— Likelihood contours (68%, 95%, 99%) in the $w - y_{D,\max}$ plane for the 21 LB LOS (top panel), the 28 nLB LOS (middle panel), and all 49 LOS (bottom panel), using the top-hat probability distribution. The dashed line is $w = y_{D,\max}$ (see Figure 2).

redshift, low-metallicity LOS to background quasars (O’Meara et al. 2006; Pettini et al. 2008), the inferred primordial D abundance, $y_{\text{DP}} = 2.8 \pm 0.2$, is in excellent agreement with the non-BBN inferred baryon density parameter (Steigman 2007). In the post-BBN Universe deuterium is destroyed as gas is cycled through stars, so that comparing the abundance of deuterium in the ISM of the Galaxy with the primordial D abundance provides an estimate of the virgin fraction of the ISM (*i.e.*, the amount of gas presently in the ISM which has never been cycled through stars), constraining models of Galactic chemical

Table 1: Results for the different shapes of the adopted priors.

Bias Shape	$y_{D,\max}$	w	f_D	$\Delta \ln \mathcal{L}_{\max}$
Top-hat	2.0 ± 0.1	1.3 ± 0.2	1.4 ± 0.1	0.5
Positive	1.9 ± 0.1	1.4 ± 0.2	1.5 ± 0.1	0.9
Negative	2.4 ± 0.2	$1.7^{+0.3}_{-0.2}$	1.2 ± 0.1	2.7
M-Shaped	1.8 ± 0.1	1.0 ± 0.1	1.6 ± 0.1	0.0
Λ -Shaped	$2.3^{+0.2}_{-0.1}$	$1.8^{+0.3}_{-0.2}$	1.2 ± 0.1	1.4




Fig. 5.— Likelihood contours (68%, 95%, 99%) in the $w - y_{D,\max}$ plane for the 21 LB LOS (top panel), the 28 nLB LOS (middle panel), and all 49 LOS (bottom panel), using the M-shaped probability distribution. The dashed line is $w = y_{D,\max}$ (see Figure 2).

evolution (Steigman & Tosi 1992, 1995; Steigman et al. 2007; Prodanović & Fields 2008). According to conventional wisdom the deuterium-free, metal-enhanced products of stellar nucleosynthesis should be well-mixed in the local ISM. In contrast, the FUSE data on the abundances of deuterium and several metals (*e.g.*, iron, oxygen, etc.) along LOS within $\sim 1 - 2$ kpc of the Sun reveal a much different picture. The FUSE (Linsky et al. 2006) and earlier observations (Jenkins et al. 1999; Sonneborn et al. 2000; Hébrard et al. 2002; Hoopes et al. 2003; Prochaska et al. 2005) reveal unexpectedly large gas phase variations in y_D (and in the abundances of iron, oxygen, etc.) within the local ISM, as shown for FUSE deuterium data in Figure 1. It has been proposed that the large variations observed in the local ISM D abundances can be accounted for by preferential depletion of deuterium (relative to hydrogen) onto dust (Jura 1982; Draine 2004, 2006), although incompletely mixed infall of relatively unprocessed, deuterium-enhanced, metal-free material may have contributed to some of the observed variations (Steigman & Tosi 1992, 1995; Steigman et al. 2007). The large variations among the ISM D abundances, along with observational errors and the possible contributions from dust depletion and infall, complicate using the D observations to provide a robust estimate of the ISM D abundance which, in combination with the primordial

D value, can lead to a constraint on the deuterium astration factor, f_D . The key question is, given the data (with its errors), how to find the best estimate of the “true”, undepleted, ISM D abundance?

Here, to address this question, the limits to the true, undepleted, ISM D abundance were investigated employing a model-independent Bayesian statistical analysis similar to that used by Hogan et al. (1997) to infer the primordial helium abundance from a set of helium abundance observations. It was assumed, along with Linsky et al. (2006), that the spread in the observed D abundances is the result of incompletely homogenized D depletion onto dust in the local ISM. In our analysis this is modeled by five different probability distributions (priors) for the y_D values. The y_D (actually, $\log y_D$) values shown in Figure 1 suggest that, given the relatively large errors, a uniform (top-hat) distribution, favoring neither low-D nor high-D may be a good approximation to the data. To explore the sensitivity of our result to the choice of the prior, we first considered two asymmetric distributions – a positive-bias prior favoring low depletion, and a negative-bias prior favoring large depletion, as well as two other priors – an M-shaped distribution favoring both low and high depletion, and a complementary, Λ -shaped distribution.

Using the FUSE deuterium observations along all 49 LOS (Linsky et al. 2006; Oliveira & Hébrard 2006; Dupuis et al. 2009), we found the likelihoods in the $\{y_{D,\max}, w\}$ plane for the five choices of the Bayesian priors (see Figures 3 and 5). For all priors, the Bayesian analysis of the full data set requires significant depletion (*e.g.*, $w \neq 0$ at greater than 99.9% confidence). Comparing the maximum likelihood values for the five different distributions, we find that the bimodal, M-shaped distribution provides the best fit to the observed data (see Table 1). However, it is important to notice that the shapes of the priors require an additional assumption in our analysis, so that the M-shaped distribution is the most model-dependent.

3

In contrast, the top-hat prior is the least model-dependent, favoring all levels of depletion equally. Given our ignorance of the detailed depletion mechanisms responsible for the observed scatter in the gas phase ISM deuterium abundances, we prefer to adopt for our estimate of the undepleted, ISM deuterium abundance, the result of the simplest, top-hat

³The M-shaped prior favors both low and high levels of D depletion while strongly disfavoring intermediate depletion, suggesting that two competing processes may be at work: depletion onto dust and evaporation from dust, perhaps due to exposure to shocks. To fit the M-prior scenario, both processes would have to be efficient and rapid to account for the deficit of intermediate D abundances. The distribution of the presently available data (with its errors) is inconclusive and does not strongly favor any of the adopted prior distributions. When more data become available, the Bayesian approach presented here may be used to learn more about the mechanism of deuterium depletion onto dust.

prior, whose maximum likelihood is similar to that of the best-fitting M-distribution,

$$y_{\text{D,ISM}} \geq y_{\text{D,max}} = 2.0 \pm 0.1 = 2.0(1 \pm 0.05) \quad (10)$$

This value is our best estimate of the true ISM D abundance based on the available deuterium observations in the local ISM and is independent of any model-dependent assumptions about galactic chemical evolution. Combining our result with $y_{\text{DP}} = 2.8 \pm 0.2 = 2.8(1 \pm 0.07)$ (Pettini et al. 2008) (which, recall, provides a *lower* bound to the primordial abundance), yields a limit to the deuterium astration factor $f_{\text{D}} \leq 1.4 \pm 0.1$ (for the M-prior, $f_{\text{D}} \leq 1.6 \pm 0.1$), consistent with most, but not all, Galactic chemical evolution models (Steigman et al. 2007; Prodanović & Fields 2008; Romano 2009). If, on the other hand we compared this $y_{\text{D,ISM}}$ value to the BBN + WMAP inferred primordial D abundance, *e.g.*, $y_{\text{DP}} = 2.5 \pm 0.1$ (Steigman 2010), $y_{\text{DP}} = 2.5 \pm 0.2$ (Cyburt et al. 2008) or, the prediction inferred when including the WMAP-determined effective number of neutrino species (Komatsu et al. 2010), $y_{\text{DP}} = 3.0 \pm 0.4$, the resulting deuterium astration factor would be somewhat lower in first two cases, $f_{\text{D}} \approx 1.3 \pm 0.1$, which is marginally problematic for some GCE models. In contrast, for the Komatsu et al. (2010) value of y_{DP} , $f_{\text{D}} \leq 1.5 \pm 0.2$, which is entirely consistent with GCE models.

As seen in Figures 1 – 4, for the LB there is little scatter among the gas-phase D abundances. The small scatter is entirely consistent with the observational errors ($w = 0$) and all LB D abundances are consistent, within the errors, with $y_{\text{D,LB}} = 1.5(1 \pm 0.03)$. This suggests that for the Local Bubble, $y_{\text{D,ISM}} \geq y_{\text{D,LB}}$ and, $f_{\text{D,LB}} \leq 1.8 \pm 0.1$, consistent with all the successful chemical evolution models identified in SRT. However, while the uniform LB D abundance suggests that D may be *undepleted* in the LB, for all LOS, $y_{\text{D,max}} \approx 1.3 y_{\text{D,LB}}$, suggesting either that D is *depleted uniformly* in the LB or, that outside of the LB the gas phase deuterium abundance may have been *enhanced* along some LOS by the addition of nearly primordial gas which has recently fallen into the disk of the Galaxy in the form of cloudlets which take some time to mix with the pre-existing gas in the ISM. Does $y_{\text{D,LB}} = 1.5$ or $y_{\text{D,max}} = 2.0$ provide the best estimate of the lower bound to the ISM D abundance? If deuterium is depleted onto dust, why is there not a strong correlation between deuterium abundance and iron depletion⁴ (Linsky et al. 2006) and, which refractory element is then best to use as proxy for determining deuterium depletion onto dust? These questions cannot

⁴As pointed out by the Referee, shock strength may play a role in accounting for the scatter observed in the correlation between the gas phase D and Fe abundances. If deuterium is loosely bound to the grain mantle while iron is locked into the core of the dust grain, deuterium would be more easily returned to the gas than iron when grains are exposed to shocks of modest strength, while iron might be removed from dust grains only by stronger shocks. The scatter in the correlation between the gas phase D and Fe abundances may be an indicator of shock strength.

be answered by the analysis presented here. In a companion paper (Steigman & Prodanović 2010), abundances of refractory elements are used in concert with the deuterium abundances in an attempt to resolve this question.

Acknowledgments

We are grateful to the Referee (J. L. Linsky) for a constructive and valuable report which has helped us improve on our original manuscript. GS acknowledges valuable discussions with D. Romano and M. Tosi and we thank M. Tosi, C. Hogan, and V. Pavlidou for helpful remarks on an earlier version of this manuscript. The work of TP is supported by the Provincial Secretariat for Science and Technological Development, and by the Ministry of Science of the Republic of Serbia under project number 141002B. The research of GS is supported at The Ohio State University by a grant from the US Department of Energy. Some of the work reported here was carried out when GS was a Humboldt Awardee at the Max Planck Institute for Physics and the Ludwig Maximilians University in Munich. GS is grateful to the AvH for its support and to the MPI and the LMU for hospitality.

REFERENCES

- Boesgaard A. M., Steigman G., 1985, *Ann. Rev. Astron. Astrophys.*, 23, 319
- Coc A., Vangioni-Flam E., Descouvemont P., Adahchour A., Angulo C., 2004, *ApJ*, 600, 544
- Cyburt R. H., Fields B. D., Olive K. A., 2003, *Phys. Lett. B*, 567, 227
- Cyburt R. H., Fields B. D., Olive K. A., 2008, *JCAP*, 11, 12
- Draine B. T., Salpeter E. E., 1979, *ApJ*, 231, 438
- Draine B. T., 2004, *Origin and Evolution of the Elements*, McWilliam A., Rauch M., eds., Cambridge Univ. Press, Cambridge, p. 317
- Draine B. T., 2006, *Astrophysics in the Far Ultraviolet: Five Years of Discovery with FUSE*, 348, 58
- Dring A. R., Linsky J., Murthy J., Henry R. C., Moos W., Vidal-Madjar A., Audouze J., Landsman W., 1997, *ApJ*, 488, 760
- Dunkley, J., et al., 2009, *ApJS*, 180, 306

- Dupuis J., Oliveira C. M., Hébrard G. H., Moos H. W., Sonnentrucker P., 2009, *ApJ*, 690, 1045
- Epstein R. I., Lattimer J. M., Schramm D. N., 1976, *Nature*, 263, 198
- Hébrard G., et al., 2002, *ApJS*, 140, 103
- Hogan C. J., Olive K. A., Scully S. T., 1997, *ApJL*, 489, L119
- Hoopes C. G., Sembach K. R., Hébrard G., Moos H. W., Knauth D. C., 2003, *ApJ*, 586, 1094
- Jenkins E. B., Tripp T. M., Woźniak P. R., Sofia U. J., Sonneborn G., 1999, *ApJ*, 520, 182
- Jura M., 1982, *Advances in Ultraviolet Astronomy*, Kondo Y., Mead J., Chapman R. D., eds., NASA, Washington, p. 54
- Komatsu, E., et al., 2009, *ApJS*, 180, 330.
- Komatsu, E., et al., (arXiv:1001:4538) [astro-ph].
- Lallement R., Welsh B. Y., Vergely J. L., Crifo F., Sfeir D., 2003, *A&A*, 411, 447
- Lallement R., Hébrard G., Welsh B. Y., 2008, *A&A*, 481, 381
- Linsky J. L., et al., 2006, *ApJ*, 647, 1106
- O’Meara J. M., Burles S., Prochaska J. X., Prochter G. E., Bernstein R. A., Burgess K. M., 2006, *ApJL*, 649, L61
- Oliveira C. M., Hébrard G., 2006, *ApJ*, 653, 345
- Pettini M., Zych B. J., Murphy M. T., Lewis A., Steidel C. C., 2008, *MNRAS*, 391, 1499
- Piskunov N., Wood B. E., Linsky J. L., Dempsey R. C., Ayres T. R., 1997, *ApJ*, 474, 315
- Prochaska J. X., Tripp T. M., Howk J. C., 2005, *ApJ*, 620, L39
- Prodanović T., Fields B. D., 2003, *ApJ*, 597, 48
- Prodanović T., Fields B. D., 2008, *JCAP*, 9, 3
- Redfield S., Linsky J. L., 2008, *ApJ*, 673, 283
- Romano D., Tosi M., Chiappini C., Matteucci F., 2006, *MNRAS*, 369, 295

- Romano, D. 2009, arXiv:0912.3737 [astro-ph]
- Schramm D. N., Turner M. S., 1998, Rev. Mod. Phys., 70, 303
- Simha V., Steigman G., 2008, JCAP, 08, 011
- Sonneborn G., Tripp T. M., Ferlet R., Jenkins E. B., Sofia U. J., Vidal-Madjar A., Woźniak P. R., 2000, ApJ, 545, 277
- Spergel D. N., et al., 2007, ApJS, 170, 377
- Steigman G., 2007, Ann. Rev. Nucl. Part. Sci, 57, 463
- Steigman G., Tosi M., 1992, ApJ, 401, 150
- Steigman G., Tosi M., 1995, ApJ, 453, 173
- Steigman G., Romano D., Tosi M., 2007, MNRAS, 378, 576
- Steigman G. 2010, arXiv:1002.3604 [astro-ph]
- Steigman G., Prodanović T., 2010, *in preparation*
- Tielens A. G. G. M. 1983, A&A, 119, 177
- Tosi M., 1996, From Stars to Galaxies: The Impact of Stellar Physics on Galaxy Evolution, 98, 299
- Vangioni-Flam E., Olive K. A., Prantzos N., 1994, ApJ, 427, 618
- Yang J., Turner M. S., Steigman G., Schramm D. N., Olive K. A., 1984, ApJ, 281, 493